

Cooling Dynamics of Ultracold Two-Species Fermi-Bose Mixtures

Carlo Presilla^{1,2,3} and Roberto Onofrio^{4,2,5}

¹*Dipartimento di Fisica, Università di Roma "La Sapienza," Piazzale Aldo Moro 2, Roma 00185, Italy*

²*Istituto Nazionale per la Fisica della Materia, Unità di Roma 1 and Center for Statistical Mechanics and Complexity, Roma 00185, Italy*

³*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, Roma 00185, Italy*

⁴*Dipartimento di Fisica "G. Galilei," Università di Padova, Via Marzolo 8, Padova 35131, Italy*

⁵*Los Alamos National Laboratory, Los Alamos, New Mexico 87545*

(Received 23 October 2002; published 24 January 2003)

We compare strategies for evaporative and sympathetic cooling of two-species Fermi-Bose mixtures in single-color and two-color optical dipole traps. We show that in the latter case a large heat capacity of the bosonic species can be maintained during the entire cooling process. This could allow one to efficiently achieve a deep Fermi degeneracy regime having at the same time a significant thermal fraction for the Bose gas, crucial for a precise thermometry of the mixture. Two possible signatures of a superfluid phase transition for the Fermi species are discussed.

DOI: 10.1103/PhysRevLett.90.030404

PACS numbers: 05.30.Fk, 32.80.Pj, 67.57.-z, 67.60.-g

Recent studies of ultracold dilute matter are bridging a gap between the idealized descriptions of quantum degenerate Bose and Fermi gases and their actual counterpart in strongly interacting condensed matter systems like liquid ⁴He and electrons in superconducting materials [1]. While experimental studies of interacting dilute Bose gases in the degenerate regime are ongoing since 1995 [2], Fermi gases have been explored only more recently. Noninteracting, purely quantum-mechanical features of dilute Fermi gases have been observed in the degenerate regime, namely, Pauli blocking [3] and Fermi pressure [4,5]. Phenomena involving their interacting nature are expected when fermions are highly degenerate. Important studies of strongly interacting degenerate Fermi gases have been recently reported for Fermi-Bose mixtures [6], and for two-component Fermi gases [7]. Moreover, BCS-based models are predicting a superfluid phase based on Cooper pairing already invoked for the understanding of low-temperature superconductivity and superfluidity of ³He [8].

Current efforts to cool fermions seem ultimately limited by intrinsic heating sources [9] in the case of evaporative cooling of two hyperfine states of fermions [3,10], and by the decreasing cooling efficiency of bosons in experiments using sympathetic cooling [4,5,11,12]. Recently, we proposed the use of a two-color optical dipole trap to enhance the Fermi degeneracy temperature T_F with respect to the Bose-Einstein critical temperature T_c whenever a bosonic species is used to sympathetically cool a different fermionic species [13]. This can be obtained by engineering different trapping potentials for the two species with proper detuning and intensities of two laser beams, as discussed in [13] with particular regard to the static confinement features. In this Letter we analyze the dynamics of cooling and heating of the Fermi-Bose mixture in two-color optical dipole traps. It turns out that a deep Fermi degeneracy regime can be achieved, allow-

ing at the same time for both precision thermometry and relatively simple signatures of fermion superfluidity.

In order to understand the efficiency limits of sympathetic cooling of a fermion-boson mixture let us consider the heat capacities $C(N, T)$ of the two species at a fixed number N of particles as a function of temperature T . In the case of noninteracting gases confined into a harmonic potential $V(x, y, z) = \frac{1}{2}m(\omega_x^2x^2 + \omega_y^2y^2 + \omega_z^2z^2)$, the heat capacities can be evaluated numerically. In Fig. 1 we show the behavior of $C_b(N_b, T)$ and $C_f(N_f, T)$ for a mixture composed by the same number of bosons and fermions trapped in a crossed-beam optical dipole trap [14,15]. Below the critical, $T_c = \zeta(3)^{-1/3}\hbar\omega_b N_b^{1/3}k_B^{-1}$, and Fermi, $T_F = 6^{1/3}\hbar\omega_f N_f^{1/3}k_B^{-1}$, temperatures, defined in




FIG. 1 (color online). Heat exchange for harmonically trapped Bose-Fermi mixtures. The single particle heat capacity of noninteracting fermions (dot-dashed) and bosons (solid) is shown versus temperature for two different values of the trap frequency ratio ω_f/ω_b . Arrows evidence the different T/T_F values below which the boson heat capacity becomes smaller than the fermion one. We consider a case with $N_b = N_f = 10^6$, $m_b = m_f$, and $\omega_x = \omega_y = \omega_z/\sqrt{2}$.

terms of the average angular trap frequencies $\omega_b = (\omega_{bx}\omega_{by}\omega_{bz})^{1/3}$ and $\omega_f = (\omega_{fx}\omega_{fy}\omega_{fz})^{1/3}$, the boson and fermion heat capacities vanish as T^3 and T , respectively. If $\omega_f = \omega_b$, the boson heat capacity becomes smaller than the fermion one below $T/T_F \approx 0.3$, strongly affecting the efficiency of sympathetic cooling for smaller T/T_F . This explains qualitatively the difficulty in reaching temperatures lower than $T/T_F \approx 0.25$ in the experiments reported in [4,5] where ${}^7\text{Li}$ - ${}^6\text{Li}$ mixtures were used, and more in general in magnetically or single-color optically trapped Fermi-Bose mixtures. Were we able to increase the ratio ω_f/ω_b , the cooling efficiency of a boson-fermion mixture could be extended to much lower temperatures. As an example, in Fig. 1 we show that for $\omega_f/\omega_b = 10$, obtainable with bichromatic optical dipole traps [13], the heat capacity inversion takes place at $T/T_F \approx 10^{-2}$. The discussion can be made more quantitative by considering the dynamics of the system during forced evaporation and comparing a bichromatic trap to the single-color case.

In optical dipole traps forced evaporative cooling is obtained by continuously decreasing the depth of the confining potential energy via proper control of the laser power. The detailed dynamics for a single-color optical dipole trap has been discussed in [16], where scaling laws for all the relevant parameters of evaporative cooling were obtained. A fundamental quantity in forced evaporation is the ratio between the potential energy depth experienced by the trapped atoms and their temperature, $\Delta U/k_B T \equiv \eta$. It has been shown that thermodynamic equilibrium is assured if η is kept constant, even if ΔU and T are time dependent [17]. This implies that during cooling the temperature of the atomic cloud is always well defined. The condition of constant η determines the time dependence of the potential energy depth [16]

$$\frac{\Delta U(t)}{\Delta U_i} = \left(1 + \frac{t}{\tau}\right)^{\varepsilon_U}, \quad (1)$$

where ΔU_i is the initial potential depth, $\varepsilon_U = -2(\eta' - 3)/\eta'$, and $\tau^{-1} = (2/3)\eta'(\eta - 4)\exp(-\eta)\gamma_i$, with $\eta' = \eta + (\eta - 5)/(\eta - 4)$ and γ_i being the initial elastic collision rate. Once the time dependence of ΔU is known, all other relevant quantities, e.g., number of particles, temperature, phase space density, and elastic scattering rate, are obtained by scaling laws similar to (1) with possibly different exponents, ε_N , ε_T , ε_ρ , and ε_γ . In Table I we report the values of these exponents for three different values of η realistically achievable in experimental situations, and the corresponding time constant τ .

In the case of a mixture of bosonic and fermionic species, Eq. (1) describes the potential energy depth of bosons ΔU_b [18]. This quantity, in turns, fixes the laser power P necessary to create the confining potential well. The potential energy depth of fermions ΔU_f is then determined as a function of P . By using the scaling law (1), we have studied evaporative cooling strategies for single-color and two-color optical dipole traps with

TABLE I. Evaporative cooling scaling exponents for the potential energy depth ε_U , the temperature ε_T , the number of particles ε_N , the phase-space density ε_ρ , and the elastic collision rate ε_γ , for three values of the evaporation parameter η . We also report the time constant τ in terms of the initial elastic scattering rate $\gamma_i = N_b m_b \sigma \omega_b^3 / (2\pi^2 k_B T)$, where N_b , ω_b , and T are the initial values of the number of bosons, their average angular trap frequency and temperature, respectively, and σ the elastic cross section.

| η | $\varepsilon_U, \varepsilon_T$ | ε_N | ε_ρ | ε_γ | $(\gamma_i \tau)^{-1}$ |
|--------|--------------------------------|-----------------|--------------------|----------------------|------------------------|
| 5 | -0.80 | -0.60 | 0.60 | -1 | 2.2×10^{-2} |
| 10 | -1.45 | -0.28 | 1.89 | -1 | 2.0×10^{-3} |
| 15 | -1.62 | -0.19 | 2.25 | -1 | 3.6×10^{-5} |

${}^6\text{Li}$ - ${}^{23}\text{Na}$ mixtures, recently brought to quantum degeneracy for both species in a magnetic trap [11]. Analogous considerations hold for all the combinations of the only two available stable Fermi alkali isotopes, ${}^6\text{Li}$ and ${}^{40}\text{K}$, refrigerated through the largely available Bose coolers, ${}^{23}\text{Na}$ and ${}^{87}\text{Rb}$. In the single-color case only a red-detuned laser beam is present and its power P_1 is decreased continuously as shown by the dashed line in the upper panel of Fig. 2. The ratio of the fermionic and bosonic average trapping frequencies has the constant value $\omega_f/\omega_b = 1.96$ determined by the masses of the two species and the different detunings of the atomic transition wavelengths with respect to the red-detuned laser wavelength. In the two-color situation a coaxial blue-detuned beam focused on the center of the existing optical dipole trap is turned on at time t_0 and for $t \geq t_0$ its power P_2 is maintained at a constant ratio with the red-detuned laser power, $P_2/P_1 = \text{const}$. By demanding a smooth time dependence for $\Delta U_b(t)$ as described by Eq. (1)—see central panel of Fig. 2—together with the fact that $\Delta U_b = \Delta U_b(P_1, P_2)$, a discontinuity at $t = t_0$ also for the red-detuned laser power is required. For $t \geq t_0$ the ratio ω_f/ω_b can be ideally increased to an arbitrary high value by choosing a proper ratio P_2/P_1 . The abrupt increase (decrease) of the fermionic (bosonic) average trapping frequency due to the turning on of the blue-detuned laser determines a corresponding decrease (increase) of the Fermi (critical) temperature. As shown in the lower panel of Fig. 3, this produces an increase of T/T_c and a decrease of T/T_F with respect to their corresponding smooth evolutions in the case of a single-color trap. Therefore, the presence of the blue-detuned laser helps to both maintain the Bose gas in a noncondensed state and allow for a deeper degeneracy condition of the Fermi gas. At the same time, evaporative cooling is less efficient as a consequence of the weakening of the confinement caused by the blue-detuned beam. However, this is not an issue since in the latest stage of evaporation we estimate elastic scattering rates of order $\Gamma_{el} \approx 10^2 \text{ s}^{-1}$, even taking into account the suppression of scattering induced by Pauli blocking [19]. Also, the blue-detuned beam gives the dominant contribution to the heating of the mixture




FIG. 2 (color online). Evaporative cooling strategies for an optically trapped ${}^6\text{Li}\text{-}{}^{23}\text{Na}$ mixture. Time evolution of red-detuned and blue-detuned laser powers (upper panel), fermion and boson trap depths (central panel), and fermion and boson trap frequencies (lower panel) for a single-color optical dipole trap (dashed lines) and a two-color optical dipole trap (solid lines). The laser powers are fixed by the condition that $\Delta U_b(t)$ follows Eq. (1) with $\eta = 10$ and $P_2/P_1 = 0$ in the single-color case or $P_2/P_1 = 0.32$ (corresponding to $\omega_f/\omega_b = 13.15$) for $\gamma_{it} \geq 400$ in the two-color case. The wavelengths of the laser beams are chosen at $\lambda_1 = 1064$ nm and $\lambda_2 = 532$ nm.

due to Rayleigh scattering of the sodium atoms, but this is largely compensated by the cooling power of the Bose gas even in the latest stage of the evaporation. Other sources of heating, like hole heating [9] or technical laser noise [20], can be made negligible with respect to the heating induced by Rayleigh scattering. In particular, fluctuations in the laser power ratio P_2/P_1 could induce instabilities and parametric heating especially in the interesting regime where the ratio ω_f/ω_b is made large. For $\omega_f/\omega_b = 10$, a power ratio stability of 0.1%–1% is required, which is within the capability of current laser stabilization techniques.

In the upper panel of Fig. 3 we show the evolution of the ratio C_b/C_f between the heat capacities of the bosonic and fermionic species—a figure of merit of the sympathetic cooling efficiency. The heat capacities have been evaluated numerically taking into account the time evolution of the relevant quantities, in particular, the diminishing number of bosons during forced evaporation which strongly affects the time dependence of C_b (while the effect of many-body interactions, evaluated in [21], gives




FIG. 3 (color online). Efficiency of sympathetic cooling in optical dipole traps. The time evolution of heat capacity ratio C_b/C_f (upper panel) and temperature ratios T/T_F and T/T_c (lower panel) are depicted for a ${}^6\text{Li}\text{-}{}^{23}\text{Na}$ mixture in which initially $N_f = 10^5$ and $N_b = 10^6$. The single-color (dashed lines) and two-color (solid lines) cases refer to the trapping configurations defined in Fig. 2. Unlike the latter, for the single-color case the heat capacity ratio approaches unity in the latest stage, and the equilibrium temperature of the mixture is no longer dominated by the bosonic component undergoing forced evaporation.

a much weaker time dependence). The boson heat capacity in the single-color trap becomes smaller than the fermion one at times $\gamma_{it} \geq 6200$ when most of the bosons are condensed ($T/T_c \approx 0.25$, $T/T_F \approx 0.1$). On the other hand, in the two-color trap C_b/C_f maintains values much larger than unity, and it is possible to reach $T/T_F \approx 0.02$ while $T/T_c \geq 0.3$. The presence of a larger bosonic thermal cloud for the two-color case even at the latest stage of fermion cooling allows for a more precise thermometry. The estimate of the temperature for the Fermi-Bose mixture is indeed obtained by fitting the tail of the normal Bose component superimposed to the condensate fraction. As the temperature is lowered, the thermal component shrinks in amplitude and size, therefore lowering the accuracy of the measurement. This effect is mitigated in the two-color trap.

Superfluidity of the Fermi gas is expected below the critical temperature for the onset of atomic Cooper pairs [8] $T_{\text{BCS}} \approx 5/3 \exp(-\pi/2k_F|a| - 1) T_F$, where k_F is the Fermi wave vector and a the elastic scattering length of fermions. Besides leaving freedom to apply arbitrary homogeneous magnetic fields to enhance the scattering length through tuning to a Feshbach resonance [22–24], our bichromatic configuration allows also for an

independent increase of k_F due to the higher achievable densities. The resulting T_{BCS}/T_F are within the explorable range which corresponds, as seen in the lower panel of Fig. 3, to $T/T_F \geq 2 \times 10^{-2}$. The presence of a superfluid state could be evidenced by using the same blue-detuned beam used to deconfine the bosons as a mechanical stirrer for the fermion cloud. Thus, in analogy to already performed experiments on Bose condensates, one could look at a finite threshold for the onset of a highly dissipative regime [25] or of a drag force [26]. The stirring of the Fermi gas occurs in the presence of both a bosonic thermal cloud and a Bose-condensed component. These last give rise to heating at all stirring velocities [27] and at a critical velocity lower than for the one expected for the Fermi superfluid, respectively. However, due to their low density in the latest cooling stage, the contributions to the heating induced by the stirring of the Bose components are much smaller than the Rayleigh heating. To discriminate against the bosonic cloud background one could take advantage of the recently proposed manipulations of an ultracold cloud with Raman beams by creating a directional critical velocity for the superfluid Fermi component [28]. An alternative signature for superfluidity consists in looking at the bulge in the density profile predicted below T_{BCS} [29]. Here, again, the presence of a thermal cloud for the bosons makes this background simpler to discriminate against any fermion superfluidity signature due to the well-controllable Gaussian-shaped profile of the former, its weaker interactions with the Fermi gas [30], and the broader Thomas-Fermi profile of the Bose-condensed component caused by the shallower confinement.

In conclusion, our analysis of evaporative and sympathetic cooling in a two-color optical dipole trap shows that a deep Fermi degeneracy regime can be achieved by efficiently exploiting the cooling capability of a Bose gas with large heat capacity. The fact that the cooler need not be in the condensed phase gives also larger flexibility for choosing the Bose species. One could reconsider the use of ^{133}Cs which, due to its large mass and small recoil temperature, can be efficiently cooled to very low temperatures by purely optical means, therefore ensuring robust conditions in terms of temperature and heat capacity to start evaporative cooling [31]. More generally, the presence of a larger thermal component for the Bose gas does not interfere significantly with the Fermi component; rather it allows for a more accurate thermometry and a more controllable background against possible signatures of the fermion superfluid phase.

We thank S. Gupta, Z. Hadzibabic, and L. Viola for useful discussions. This work was supported in part by Cofinanziamento MIUR protocollo MM02263577_001.

[1] C. J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases* (Cambridge University Press, Cambridge, 2002).

- [2] M. H. Anderson *et al.*, *Science* **269**, 198 (1995); K. B. Davis *et al.*, *Phys. Rev. Lett.* **75**, 3969 (1995); C. C. Bradley, C. A. Sackett, J. J. Tollett, and R. G. Hulet, *Phys. Rev. Lett.* **75**, 1687 (1995); **79**, 1170 (1997).
- [3] B. DeMarco and D. S. Jin, *Science* **285**, 1703 (1999).
- [4] A. G. Truscott *et al.*, *Science* **291**, 2570 (2001).
- [5] F. Schreck *et al.*, *Phys. Rev. Lett.* **87**, 080403 (2001).
- [6] G. Modugno *et al.*, *Science* **297**, 2240 (2002).
- [7] K. M. O'Hara *et al.*, *Science* **298**, 2179 (2002).
- [8] H. T. C. Stoof, M. Houbiers, C. A. Sackett, and R. G. Hulet, *Phys. Rev. Lett.* **76**, 10 (1996).
- [9] E. Timmermans, *Phys. Rev. Lett.* **87**, 240403 (2001). The heating source discussed in this paper should not be a practical limitation if a background pressure in the trapping region $< 10^{-11}$ Torr is achieved, a requirement already fulfilled in various apparatuses.
- [10] S. R. Granade *et al.*, *Phys. Rev. Lett.* **88**, 120405 (2002).
- [11] Z. Hadzibabic *et al.*, *Phys. Rev. Lett.* **88**, 160401 (2002).
- [12] G. Roati *et al.*, *Phys. Rev. Lett.* **89**, 150403 (2002).
- [13] R. Onofrio and C. Presilla, *Phys. Rev. Lett.* **89**, 100401 (2002).
- [14] C. S. Adams *et al.*, *Phys. Rev. Lett.* **74**, 3577 (1995).
- [15] M. D. Barrett, J. A. Sauer, and M. S. Chapman, *Phys. Rev. Lett.* **87**, 010404 (2001).
- [16] K. M. O'Hara *et al.*, *Phys. Rev. A* **64**, 051403R (2001).
- [17] W. Ketterle and N. J. Van Druten, in *Advances in Atomic, Molecular, and Optical Physics*, edited by B. Bederson and H. Walther (Academic Press, San Diego, 1996), Vol. 37, p. 181.
- [18] We assume that the presence of the Fermi gas does not affect significantly evaporative cooling of the Bose gas. This requires that the potential energy depth of fermions ΔU_f is larger than that of bosons ΔU_b , and that $C_b/C_f \gg 1$.
- [19] M. J. Holland, B. DeMarco, and D. S. Jin, *Phys. Rev. A* **61**, 053610 (2000).
- [20] M. E. Gehm *et al.*, *Phys. Rev. A* **58**, 3914 (1998).
- [21] S. Giorgini, L. P. Pitaevskii, and S. Stringari, *J. Low Temp. Phys.* **109**, 309 (1997).
- [22] E. Timmermans *et al.*, *Phys. Lett. A* **285**, 228 (2001).
- [23] M. Holland, S. J. J. M. F. Kokkelmans, M. L. Chiofalo, and R. Walser, *Phys. Rev. Lett.* **87**, 120406 (2001).
- [24] Y. Ohashi and A. Griffin, *Phys. Rev. Lett.* **89**, 130402 (2002).
- [25] C. Raman *et al.*, *Phys. Rev. Lett.* **83**, 2502 (1999).
- [26] R. Onofrio *et al.*, *Phys. Rev. Lett.* **85**, 2228 (2000).
- [27] C. Raman *et al.*, *J. Low Temp. Phys.* **122**, 99 (2001).
- [28] J. Higbie and D. M. Stamper-Kurn, *Phys. Rev. Lett.* **88**, 090401 (2002).
- [29] M. L. Chiofalo *et al.*, *Phys. Rev. Lett.* **88**, 090402 (2002).
- [30] M. Amoruso *et al.*, *Eur. Phys. J. D* **4**, 261 (1998).
- [31] Recently, sympathetic cooling of ^7Li through ^{133}Cs has been demonstrated in a far-off resonance optical trap; see M. Mudrich *et al.*, *Phys. Rev. Lett.* **88**, 253001 (2002). In this experiment the optical potential was created by a CO_2 laser obtaining a deeper confinement for the latter species and limitations in the sympathetic cooling of the former one. This problem could be circumvented with a two-color optical dipole trap based on the use of a Nd:YAG laser as the red-detuned laser, and a deconfining beam with wavelength in between the two atomic transition wavelengths.